

C^* -SIMPLICITY OF FREE PRODUCTS WITH AMALGAMATION AND RADICAL CLASSES OF GROUPS

NIKOLAY A. IVANOV AND TRON OMLAND

ABSTRACT. We give new characterizations to ensure that a free product of groups with amalgamation has a simple reduced group C^* -algebra, and provide a concrete example of an amalgam with trivial kernel, such that its reduced group C^* -algebra has a unique tracial state, but is not simple.

Moreover, we show that there is a radical class of groups for which the reduced group C^* -algebra of any group is simple precisely when the group has a trivial radical corresponding to this class.

1. INTRODUCTION

Groups have been among the most studied objects in connection with operator algebras, and one of the natural questions to consider in this regard is whether group C^* -algebras are simple. While full group C^* -algebras can never be simple (unless the group is trivial), the problem for reduced group C^* -algebras has been of great interest ever since the work of Powers [26], showing that nonabelian free groups have a simple reduced group C^* -algebra. Larger classes of groups with properties similar to the one Powers described were later studied, and several results and open problems on this topic are discussed by de la Harpe [17].

Recall that a discrete group G is called C^* -simple if its reduced group C^* -algebra $C_r^*(G)$ is simple, and G is said to have the unique trace property if $C_r^*(G)$ has a unique tracial state. A long-standing open problem was whether the two properties were equivalent, but fairly recently it has been shown that C^* -simplicity is strictly stronger than the unique trace property. The “stronger” part is due to Breuillard, Kalantar, Kennedy, and Ozawa [8, 21] and the “strictly” part is due to Le Boudec [4]. The former showed that the unique trace property is equivalent to having a trivial amenable radical, a property already known to be weaker than C^* -simplicity, see [17]. Both of the works [8, 21] use extensively the theory of boundary actions developed by Furstenberg and Hamana. Even more recently, some other more operator-theoretical characterizations have been obtained by Haagerup [14] and Kennedy [22]. However, the conditions from [14, 22] that are equivalent with C^* -simplicity, are not always easy to check in concrete situations, for example by combinatorial group properties. We also note that the unique trace property for G always implies that G is icc, that is, every nontrivial conjugacy class in G is infinite, and it is well-known that the group von Neumann algebra associated to G is a factor if and only if G is icc.

The problem of finding conditions to ensure that a free product of groups with amalgamation is C^* -simple was first considered by Bédos [1], and mentioned as Problem 27 in [17].

In this article, we first make a detailed study of amalgamated free products, inspired by work of de la Harpe and Préaux [19]. By making use of a few new observations, we are able to improve some of the results in [19]. Then we present in Section 4 a concrete example of an amalgam that has the unique trace property, but is not C^* -simple.

Date: December 13, 2016.

2010 Mathematics Subject Classification. 46L05, 22D25 (Primary) 43A07, 20M11 (Secondary).

Key words and phrases. C^* -simplicity, free product of groups with amalgamation, radical classes of groups.

In particular, our example shows that a free product with amalgamation can fail to be C^* -simple when it has a trivial kernel, but has “one-sided” kernels that are nontrivial (and amenable). Every amalgam $G_0 *_H G_1$ acts on its Bass-Serre tree, and in this setup the one-sided kernels consist of the elements that fix all vertices of the “half-trees” obtained by removing the edge corresponding to H . This has similarities with [4].

We keep the first sections mostly to “elementary” proofs, restricting to combinatorial group properties, and delay the involvement of boundary actions to Section 5.

In the final two sections, we first recall the definitions of radical and residual classes of groups and prove several statements that will be of later use. The result of [8] implies that the class of groups with the unique trace property is the residual class “dual” to the radical class of amenable groups. We then show that the class of C^* -simple groups is also a residual class, giving rise to a “predual” radical class of groups that contains all the amenable groups, and we will call a group “amenabl-ish” if it belongs to this class. The example we provide of an amalgam that is not C^* -simple, but has the unique trace property, turns out to be a (nonamenable) amenabl-ish group. Moreover, for every group there is an amenabl-ish radical such that C^* -simplicity of the group is equivalent to this radical being trivial, giving an analog of the relation between the amenable radical and the unique trace property.

2. PRELIMINARIES

We consider only discrete groups in this article.

Let G be a group. As usual, we equip the Hilbert space $\ell^2(G)$ with the standard orthonormal basis $\{\delta_g\}_{g \in G}$, and define the left regular representation λ of G on $\ell^2(G)$ by $\lambda(g)\delta_h = \delta_{gh}$. Then the reduced group C^* -algebra of G , denoted by $C_r^*(G)$, is the C^* -subalgebra of $B(\ell^2(G))$ generated by $\lambda(G)$. The group G is called *C^* -simple* if $C_r^*(G)$ is simple, that is, if it has no nontrivial proper two-sided closed ideals.

A state on a unital C^* -algebra A is a linear functional $\phi: A \rightarrow \mathbb{C}$ that is positive, i.e., $\phi(a) \geq 0$ whenever $a \in A$ and $a \geq 0$, and unital, i.e., $\phi(1) = 1$. A state ϕ is called *tracial* if it satisfies the additional property that $\phi(ab) = \phi(ba)$ for all $a, b \in A$. There is a canonical faithful tracial state τ on $C_r^*(G)$, namely the vector state associated with δ_e , that is, $\tau(a) = \langle a\delta_e, \delta_e \rangle$ for all $a \in C_r^*(G)$. The group G is said to have the *unique trace property* if τ is the only tracial state on $C_r^*(G)$.

Recall that a group G is *amenable* if there exists a state on $\ell^\infty(G)$ which is invariant under the left translation action by G . It is explained by Day [10] that every group G has a unique maximal normal amenable subgroup, called the *amenable radical* of G . Then [8, Theorem 1.3] shows that G has the unique trace property if and only if the amenable radical of G is trivial, so C^* -simplicity is stronger than the unique trace property by [17, Proposition 3].

The first larger class of groups that were shown to be C^* -simple with the unique trace property, was the so-called Powers groups introduced by de la Harpe [16] (see also [18]).

Definition 2.1. A group G is called a *Powers group* if for any finite subset F of $G \setminus \{e\}$ and any integer $k \geq 1$ there exists a partition $G = D \sqcup E$ and elements g_1, g_2, \dots, g_k in G such that

$$\begin{aligned} fD \cap D &= \emptyset \text{ for all } f \in F, \\ g_i E \cap g_j E &= \emptyset \text{ for all distinct } 1 \leq i, j \leq k. \end{aligned}$$

Moreover, a group G is called a *weak Powers group* if it satisfies the above definition for all finite sets F that are contained in a nontrivial conjugacy class. Finally, G is called a *weak* Powers group* if it satisfies the above definition for all nontrivial one-element sets F .

Clearly, every Powers group is a weak Powers group and every weak Powers group is a weak* Powers group. It is known that the weak Powers property implies C^* -simplicity [3], while the weak* Powers property implies the unique trace property [29, Section 5.2].

Lemma 2.2. *Assume that N is a normal subgroup of a group G . If G is a Powers group, then N is a Powers group. If G is a weak Powers group, then N is a weak Powers group. If G is a weak* Powers group, then N is a weak* Powers group.*

Proof. The first statement is a (seemingly unnoticed) result by Kim [23, Theorem 1], and the last two statements are identically proven. For the convenience of the reader, we present the argument of the middle one:

Let $k \geq 1$, and suppose that F is a finite set and C' a nontrivial conjugacy class in N such that $F \subseteq C'$. If two elements are conjugate in N they are also conjugate in G , so there is a nontrivial conjugacy class C in G such that $F \subseteq C$. Since G is weak Powers, there exists a partition $G = D \sqcup E$ and g_1, g_2, \dots, g_k such that $fD \cap D = \emptyset$ for all $f \in F$ and $g_i E \cap g_j E = \emptyset$ whenever $i \neq j$.

If $k = 1$, set $D' = \{e\}$, $E' = N \setminus \{e\}$, and $s_1 = e$.

If $k \geq 2$, set $D' = D \cap N$ and $E' = E \cap N$, so $N = D' \sqcup E'$ and $fD' \cap D' \subseteq fD \cap D = \emptyset$ for all $f \in F$.

Furthermore, fix an $f \in F$ and note that the sets $g_1^{-1}g_i E$ are mutually disjoint for $i \geq 1$, so that in particular, for $i \geq 2$, then $g_1^{-1}g_i E \cap E = \emptyset$, i.e., $g_1^{-1}g_i E \subseteq G \setminus E = D$. This again implies that $f g_1^{-1}g_i E \cap D \subseteq fD \cap D = \emptyset$, so $f g_1^{-1}g_i E \subseteq G \setminus D = E$.

Set $s_1 = e$ and $s_i = g_1^{-1}g_i f g_i^{-1}g_1$ for $i \geq 2$. Since N is normal in G , we have $s_i \in N$ for all i , and if $i \neq j$, then

$$\begin{aligned} s_i E' \cap s_j E' &= g_1^{-1}g_i f g_i^{-1}g_1 E' \cap g_1^{-1}g_j f g_j^{-1}g_1 E' \subseteq g_1^{-1}g_i f g_i^{-1}g_1 E \cap g_1^{-1}g_j f g_j^{-1}g_1 E \\ &\subseteq g_1^{-1}g_i E \cap g_1^{-1}g_j E = \emptyset. \end{aligned}$$

Kim also proves that D' and E' are nonempty, but that does not seem to be necessary. \square

In [19] a group is said to be “strongly Powers” if every subnormal subgroup is a Powers group. A consequence of the above lemma is that the notion of strongly Powers coincides with Powers.

We conclude this section by mentioning two definitions. First, a group G has the *free semigroup property* if for any finite subset F of G there exists g in G such that gF is semifree, that is, the subsemigroup generated by gF in G is free over gF . Finally, a group G is said to have *stable rank one* if its reduced group C*-algebra has stable rank one, that is, the invertible elements of $C_r^*(G)$ are dense in $C_r^*(G)$.

3. FREE PRODUCTS OF GROUPS WITH AMALGAMATION

Recall that (see e.g. [28]) a free product of groups G_0 and G_1 with amalgamation over a common subgroup H (embedded via injective homomorphisms $H \rightarrow G_i$ for $i = 0, 1$) is a group G together with homomorphisms $\phi_i: G_i \rightarrow G$ for $i = 0, 1$, that agree on H , universal in the sense that for any other group G' with homomorphisms $\phi'_i: G_i \rightarrow G'$ that agree on H , there is a unique homomorphism $\phi: G \rightarrow G'$ such that $\phi'_i = \phi \circ \phi_i$.

The amalgamated free products $G = G_0 *_H G_1$ that we consider in this article are always assumed to be nondegenerate in the sense that $([G_0 : H] - 1) \cdot ([G_1 : H] - 1) \geq 2$, otherwise the situation is very different.

Let $\ker G = \bigcap_{g \in G} gHg^{-1}$ denote the kernel of the amalgam. It coincides with the normal core of H in G , i.e., the largest subgroup of H which is normal in G (and thus in G_0 and G_1).

If N is any subgroup of H which is normal in G , then the quotient G/N is isomorphic to $(G_0/N) *_H/N (G_1/N)$. In particular, this holds when $N = \ker G$. Note that we always have $\ker(G/\ker G) = \{e\}$. Indeed, if $\ker(G/\ker G) = N \subseteq H/\ker G$, let N' be the inverse image of N under the quotient map $G \rightarrow G/\ker G$. Then N' is a normal subgroup of G containing $\ker G$ and sitting inside H , so we must have $N' = \ker G$ by maximality of $\ker G$.

If G is nondegenerate, then $G/\ker G$ is nondegenerate, as $[G_i : H] = [G_i/\ker G : H/\ker G]$.

Let $FC(G)$ denote the normal subgroup of G consisting of elements with finite conjugacy class in G . Clearly, if $g \in G_i \setminus H$, then the conjugacy class of g in G must be infinite. Hence, $FC(G)$ is contained in H .

Let now $AR(G)$ denote the amenable radical of G and let $NF(G)$ denote the largest normal subgroup of G that does not contain any nonabelian free subgroup. By [9, Proposition 7], these groups fit in general into a sequence of subgroups, namely

$$(1) \quad FC(G) \subseteq AR(G) \subseteq NF(G) \subseteq \ker G \subseteq H.$$

In particular, $NF(G) \subsetneq G$, so G always contains a nonabelian free group. Moreover, there is an even longer sequence

$$FC(G) \subseteq FC(\ker G) \subseteq AR(\ker G) = AR(G) \subseteq NF(G) = NF(\ker G) \subseteq \ker G.$$

The first containment is clear, the second holds since $FC(\ker G)$ is an amenable normal subgroup of $\ker G$, and the two equalities follow from Examples 6.4, 6.6, and Lemma 6.7.

If H is finite, then $FC(G) = \ker G$, so the sequence collapses. Indeed, if $h \in \ker G$, then $h \in gHg^{-1}$ for all $g \in G$, so $g^{-1}hg \in H$ for all $g \in G$. Hence, the conjugacy class of h is contained in H , which is finite, so $h \in FC(G)$. See Theorem 3.7 for more on this case.

Proposition 3.1. *A nondegenerate amalgamated free product G has the unique trace property if and only if $\ker G$ has the unique trace property.*

Proof. Since $AR(G) = AR(\ker G)$ as explained above, this follows from [8, Theorem 1.3]. \square

Let us now recall the normal form for an element in G . First, for $i = 0, 1$, we choose sets S_i so that $S_i \cup \{e\}$ form systems of representatives for the left cosets of G_i/H . Then every element of G can be written uniquely as either $s_0 s_1 \cdots s_{n-1} s_n h$ or $s_1 \cdots s_{n-1} s_n h$, where $s_i \in S_{i \pmod{2}}$ and $h \in H$. To avoid division into separate cases, the notation $(s_0) s_1 s_2 \cdots s_{n-1} s_n h$ for the normal form of an element of G is often used, especially in Section 5.

Since we will not always assume that elements are on normal form, the following observation is sometimes useful: for $i = 0, 1$, if $g_i \in G_i \setminus H$ and $h \in H$, then there exists $g'_i \in G_i \setminus H$ such that $g_i h = h g'_i$. In other words, we can “cycle” a letter from H through a word $(g_0) g_1 \cdots g_n$ in G , with $g_i \in G_{i \pmod{2}} \setminus H$, without changing the length of the word.

Let G be any nondegenerate free product with amalgamation. We decompose G as follows. For $j = 0, 1$ and $k \geq 1$, let

$$T_{j,k} = \{g_0 \cdots g_{k-1} : g_i \in G_{i+j \pmod{2}} \setminus H\},$$

i.e., $T_{j,k}$ consists of all words of length k starting with a letter from $G_j \setminus H$. To simplify notation, we set $T_{0,0} = T_{1,0} = H$. For $j = 0, 1$ and every $k \geq 0$, we now define the set

$$(2) \quad C_{j,k} = \bigcap_{g \in T_{j,k}} g H g^{-1},$$

and note that $C_{0,0} = C_{1,0} = H$ and that $H \cap C_{j,k}$ is always a normal subgroup of H . Then

$$\ker G = \bigcap_{\substack{k \geq 0 \\ j=0,1}} C_{j,k},$$

which, as mentioned above, is always normal in G . Next, we set

$$(3) \quad K_0 = \bigcap_{k \geq 0} C_{0,k} \quad \text{and} \quad K_1 = \bigcap_{k \geq 0} C_{1,k}.$$

Both K_0 and K_1 are normal subgroups of H . Remark that it actually follows that

$$(4) \quad K_0 = \bigcap_{k \geq 0} C_{0,2k} \quad \text{and} \quad K_1 = \bigcap_{k \geq 0} C_{1,2k}.$$

Indeed, if $h \notin K_0$, then $g^{-1}hg \notin H$ for some $g \in T_{0,n}$ with $n \geq 1$. If n is odd, take any $g_1 \in G_1 \setminus H$, and note that $gg_1 \in T_{0,n+1}$ and $g_1^{-1}g^{-1}hgg_1 \notin H$, so $h \notin gg_1Hg_1^{-1}g^{-1}$, hence, $h \notin \bigcap_{k \geq 0} C_{0,2k}$. A similar argument works for K_1 .

Moreover, it should be clear that $\ker G = K_0 \cap K_1$ and that

$$(5) \quad K_0 = H \cap \bigcap_{g_0 \in G_0 \setminus H} g_0 K_1 g_0^{-1} \quad \text{and} \quad K_1 = H \cap \bigcap_{g_1 \in G_1 \setminus H} g_1 K_0 g_1^{-1}.$$

If $K_0 = \ker G$, then $K_1 \subseteq \ker G$ since $\ker G$ is normal, so $K_0 = K_1$ (similarly if $K_1 = \ker G$). Also, we have that $K_0 = \{e\}$ if and only if $K_1 = \{e\}$.

Finally, using the notation from [19, (i) p. 2-3] we set

$$(6) \quad C_k = \bigcap_{\substack{0 \leq n \leq k \\ j=0,1}} C_{j,n}.$$

Note that for any $g \in T_{0,k+1}$, we have $K_0 \subseteq gC_kg^{-1}$, and for any $g \in T_{1,k+1}$, we have $K_1 \subseteq gC_kg^{-1}$. Thus, if C_k is trivial for some $k \geq 0$, then $K_0 = K_1 = \{e\}$. Indeed, pick $k \geq 0$, $g \in T_{0,k+1}$, $h \in K_0$, and let s be an arbitrary element of length $\leq k$. Then $gs \in T_{0,n}$ for some $n \geq 1$, so $h \in gsHs^{-1}g^{-1}$. Since, this holds for all such s , we have $h \in gC_kg^{-1}$.

Moreover, it is worth noticing the difference between K_0 , K_1 , and C_k :

- the intersections defining the C_k 's involve conjugation of elements of finite length (bounded by some k), while the first letter of the words can come from any of G_0, G_1 .
- the intersections defining the K_i 's involve conjugation of elements of arbitrary length, where the first letter of the words comes from the same group.

Theorem 3.2. *Let $G = G_0 *_H G_1$ be a nondegenerate free product with amalgamation, and let K_0, K_1 be as defined above. Then the following are equivalent:*

- (i) $K_0 = K_1 = \{e\}$.
- (ii) for every finite $F \subseteq G \setminus \{e\}$, there exists $g \in G$ such that $gFg^{-1} \cap H = \emptyset$.
- (iii) for every finite $F \subseteq H \setminus \{e\}$, there exists $g \in G$ such that $gFg^{-1} \cap H = \emptyset$.

Moreover, any one of these equivalent conditions implies that G is a Powers group and has the free semigroup property.

Proof. It is obvious that (ii) implies (iii).

To see that (iii) implies (i) suppose first that $\ker G \neq \{e\}$. Then pick $f \in \ker G \setminus \{e\}$ and set $F = \{f\}$ and clearly $gfg^{-1} \in H$ for all $g \in G$, i.e., $gfg^{-1} \in gFg^{-1} \cap H$. Next, assume that $\ker G = \{e\}$, but $K_0 \neq K_1$. Recall from (5) and the subsequent note that this means both $K_0 \neq \{e\}$ and $K_1 \neq \{e\}$. Thus we pick $f_i \in K_i \setminus \{e\}$ for $i = 0, 1$ and set $F = \{f_0, f_1\}$. Let $g \in G$ be arbitrary. If $g \in H$ there is nothing to show, so assume that g ends with a letter from $G_i \setminus H$. But then $gf_i g^{-1} \in H$, i.e., $gf_i g^{-1} \in gFg^{-1} \cap H$.

Finally we prove that (i) implies (ii). Choose an arbitrary finite set $F \subseteq G \setminus \{e\}$. The idea is to conjugate the elements from F out of H one by one, and at the same time make sure we do not conjugate any elements back into H . We will repeatedly use that if $g_i \in G_i \setminus H$ and $g \notin G_i$, then $g_i g g_i^{-1} \notin H$.

Assume first there is an element $f_1 \in F \cap H$ (else there is nothing to show). Because K_0 is trivial, then (4) means that all elements in H can be conjugated out of H by an element of even length starting with a letter from $G_0 \setminus H$. That is, we can find $r_1 = g_0 g_1 \cdots g_{2n_1-1}$ such that $g_i \in G_{i \pmod{2}} \setminus H$ and $r_1^{-1} f_1 r_1 \notin H$.

Next, consider the set $F_1 = \{r_1^{-1} f r_1 : f \in F\}$. Assume that there is an element $f_2 \in F$, so that $r_1^{-1} f_2 r_1 \in H$ (otherwise we are done). Then there exists $r_2 = g'_0 g'_1 \cdots g'_{2n_2-1}$ such that $g'_i \in G_{i \pmod{2}} \setminus H$ and $r_2^{-1} r_1^{-1} f_2 r_1 r_2 \notin H$. This also means that $r_2^{-1} r_1^{-1} f_1 r_1 r_2 \notin H$.

Now we set $F_2 = \{r_2^{-1}r_1^{-1}fr_1r_2 : f \in F\}$. If all elements of F_2 are outside H we are done, so assume that there is some $f_3 \in F$ such that $r_2^{-1}r_1^{-1}f_3r_1r_2 \in H$.

It should be clear how this process continues, and since F is finite, we take r to be the product of the r_i 's, and then $r^{-1}fr \notin H$ for every $f \in F$.

The last two observations follow from [16, Proposition 10] and [13, Example 4.4 (iii),(iv), and Remark 4.5]. \square

Remark 3.3. The above result shows that [19, Theorem 3 (i)] can be slightly improved, as Lemma 2.2 shows that “strongly Powers” is the same as Powers. In fact, using the comment following (6), one can replace “ $C_k = \{e\}$ for some k ” with any of the equivalent properties of Theorem 3.2. Additionally, the countability assumption is no longer needed.

We will come back to the geometric interpretation of K_0 and K_1 in Section 5. In particular, Proposition 5.3 below gives more properties equivalent to those in Theorem 3.2.

Remark 3.4. In Section 4, we give an explicit example of a group Γ for which $\ker \Gamma = \{e\}$, while K_0 and K_1 are both nontrivial. We show that the group has the unique trace property, but is not C^* -simple. This gives a counterexample to the first author’s statement [20, Corollary 4.7], which the second author noticed to be incorrect.

Proposition 3.5. *If G is countable, H is amenable, and K_0 or K_1 is nontrivial, then G is not C^* -simple.*

This result is generalized in Theorem 5.9 below, but in the context of recent works, we provide two other short proofs of the statement.

First proof. We will show that H is recurrent in G in the sense of [22, Definition 1.4]. Let (g_n) be a sequence in G . Then at least one of the following holds:

- (a) infinitely many elements from (g_n) belongs to H
- (b) infinitely many elements from (g_n) start with a letter from $G_0 \setminus H$
- (c) infinitely many elements from (g_n) start with a letter from $G_1 \setminus H$

Pick a subsequence (g_{n_k}) of (g_n) with elements from the one of (a),(b),(c) that holds. If it is (a) then $H = \bigcap_k g_{n_k} H g_{n_k}^{-1}$, if it is (b) then $K_0 \subseteq \bigcap_k g_{n_k} H g_{n_k}^{-1}$, and if it is (c) then $K_1 \subseteq \bigcap_k g_{n_k} H g_{n_k}^{-1}$. If one of K_0 and K_1 is nontrivial, then both the other one and H are nontrivial as well. Hence, it follows from [22, Theorem 1.5] that G is not C^* -simple. \square

Second proof. If $K_0 = \ker G$, then $K_1 = \ker G$ by the comment following (5). Thus, by assumption, $\ker G$ is a nontrivial normal amenable subgroup of G , and hence G cannot be C^* -simple. The similar argument holds if $K_1 = \ker G$, so we may assume that both K_0 and K_1 are different from $\ker G$.

Choose $a \in K_0 \setminus \ker G$ and $b \in K_1 \setminus \ker G$. Then

$$\begin{aligned} \{gH : gH \neq agH\} &\subseteq \{gH : g \in T_{1,k} \text{ for some } k \geq 1\} \quad \text{and} \\ \{gH : gH \neq bgH\} &\subseteq \{gH : g \in T_{0,k} \text{ for some } k \geq 1\}, \end{aligned}$$

which are clearly disjoint. By using the technique from [15], explained in [25, p. 12], the action of G on G/H gives rise to a unitary representation $\pi : G \rightarrow \ell^2(G/H)$, that extends to a continuous representation of $C_r^*(G)$. It follows that $(1 - \lambda(a))(1 - \lambda(b))$ generates a proper two-sided closed ideal of $C_r^*(G)$. Hence, G is not C^* -simple. \square

Example 3.6. For any triple of groups H , G_0 , and G_1 , we have

$$G = (G_0 \times H) *_H (H \times G_1) \cong (G_0 * G_1) \times H.$$

In this case $H = \ker G = K_0 = K_1$, and G is C^* -simple if and only if H is C^* -simple. In particular, this means that G can be C^* -simple even if $\ker G$ is nontrivial.

We now consider the special case where H is finite.

Theorem 3.7. *Let $G = G_0 *_H G_1$ be a nondegenerate free product with amalgamation, and assume that H is finite.*

Then the following are equivalent:

- (i) G is icc
- (ii) $\ker G = \{e\}$
- (iii) $K_0 = K_1 = \{e\}$
- (iv) G is Powers
- (v) G is C^* -simple
- (vi) G has the unique trace property
- (vii) there exists $g \in G$ such that $H \cap gHg^{-1} = \{e\}$
- (viii) G has the free semigroup property

Proof. (i) \implies (ii): Since $\ker G \subseteq H$, it is a finite normal subgroup of G , so it must be trivial when G is icc.

(ii) \implies (iii): Let C_k be defined as in (6), so that $\ker G$ is the intersection of the decreasing chain $C_0 \supseteq C_1 \supseteq C_2 \supseteq \dots$. Since all the C_k 's are subgroups of H , they are finite, so if $\ker G = \{e\}$, we must have $C_k = \{e\}$ for some k . Hence, the remark following (6) explains that $K_0 = K_1 = \{e\}$.

(iii) \implies (iv) follows from Theorem 3.2.

(iv) \implies (v) \implies (vi) \implies (i) is known to hold for all groups.

(iii) \implies (vii) is also a consequence of Theorem 3.2, by taking $F = H \setminus \{e\}$.

(vii) \implies (viii) follows from [13, Proposition 5.1].

(viii) \implies (ii): If H is finite and $\ker G \neq \{e\}$, then there is no $g \in G$ such that $g \ker G$ is semifree, i.e., G does not have the free semigroup property. \square

Remark 3.8. If G is an amalgam with finite H satisfying the equivalent conditions of Theorem 3.7, then G has stable rank one by [13, Theorem 1.6] (where the argument is based on [12, Corollary 3.9] for trivial H). However, there also exists amalgams G with finite H such that G has stable rank one, but G does not satisfy the conditions of Theorem 3.7, as explained in the paragraph following [13, Theorem 1.6].

Moreover, it follows from condition (vii) and [20, Corollary 3.6] that for $G = G_0 *_H G_1$, the positive cone of the \mathbf{K}_0 -group of $C_r^*(G)$ is not perforated, i.e.,

$$\mathbf{K}_0(C_r^*(G))^+ = \{\gamma \in \mathbf{K}_0(C_r^*(G)) : \mathbf{K}_0(\tau)(\gamma) > 0\} \cup \{0\},$$

where τ is the canonical tracial state on $C_r^*(G)$. Indeed, the assumption $\mathbf{K}_1(C_r^*(H)) = 0$ is easily seen to hold from the fact that for a finite group H , $C_r^*(H)$ is a direct sum of matrix algebras and those have trivial \mathbf{K}_1 -groups.

4. AN EXAMPLE

In this section \oplus , when applied to indices, will denote summation modulo 2, and \mathbb{N} will denote the positive integers.

The group $\Gamma = G_0 *_H G_1$ is defined as follows: first H is given by the set of generators

$$\{h(i_1, i_2, \dots, i_n) : n \in \mathbb{N} \text{ and } i_k \in \{0, 1\} \text{ for all } k \in \{1, \dots, n\}\},$$

with the relations

$$h(i_1, i_2, \dots, i_n)^2 = e \text{ for all } n \in \mathbb{N}, k \in \{1, \dots, n\}, \text{ and } i_k \in \{0, 1\},$$

and for $n \geq k$

$$h(i_1, \dots, i_k)h(j_1, \dots, j_n)h(i_1, \dots, i_k) = \begin{cases} h(j_1, \dots, j_k, j_{k+1} \oplus 1, \dots, j_n) & \text{if } n > k \text{ and } i_\ell = j_\ell \text{ for all } 1 \leq \ell \leq k, \\ h(j_1, \dots, j_k, j_{k+1}, \dots, j_n) & \text{otherwise.} \end{cases}$$

Then define

$$\Gamma = \langle H \cup \{g_0, g_1\} \rangle,$$

with the additional relations

$$\begin{aligned} g_0^2 &= g_1^2 = e, & (g_0 h(1))^3 &= e, & (g_1 h(0))^3 &= e, \\ g_0 h(1, 0, i_3, \dots, i_n) g_0 &= h(0, i_3, \dots, i_n), & g_0 h(1, 1, i_3, \dots, i_n) g_0 &= h(1, 1, i_3, \dots, i_n), \\ g_1 h(0, 0, i_3, \dots, i_n) g_1 &= h(0, 0, i_3, \dots, i_n), & g_1 h(0, 1, i_3, \dots, i_n) g_1 &= h(1, i_3, \dots, i_n). \end{aligned}$$

Finally, let

$$G_0 = \langle H \cup \{g_0\} \rangle \quad \text{and} \quad G_1 = \langle H \cup \{g_1\} \rangle,$$

so that $\Gamma = G_0 *_H G_1$. Note also that

$$H = \langle \underbrace{\{h(0, \dots, 0)\}}_{n \text{ times}}, n \in \mathbb{N} \rangle \cup \langle \underbrace{\{h(1, 0, \dots, 0)\}}_{n-1 \text{ times}}, n \in \mathbb{N} \rangle.$$

Lemma 4.1. *For every $g \in G_0 \setminus H$, there exists $h \in H$ such that either $g = g_0 h$ or $g = h(1)g_0 h$, and for every $g \in G_1 \setminus H$, there exists $h \in H$ such that either $g = g_1 h$ or $g = h(0)g_1 h$.*

Consequently, $[G_0 : H] = [G_1 : H] = 3$, and the sets

$$(7) \quad S_0 = \{g_0, h(1)g_0\} \quad \text{and} \quad S_1 = \{g_1, h(0)g_1\}.$$

provide representatives for the nontrivial left cosets of G_0/H and G_1/H , respectively.

Proof. First, if h is one of the generators of H and $h \neq h(1)$, then the defining relations above show that there exists an h' such that $hg_0 = g_0 h'$. Moreover, for $h \in H$, with $h \neq h(1)$, we have $hh(1) = h(1) \cdot h(1)hh(1) = h(1)h'$ for some $h' \neq h(1)$, that is, we can move the $h(1)$'s to the left in the product. Hence, for any element $g = (g_0)h_1 g_0 h_2 g_0 \cdots g_0 h_n (g_0) \in G_0$, where $h_i \in H$, we can combine the two observations above to find the required h .

A similar argument holds in G_1 . □

Let us now denote the subgroups of H with elements determined by a prescribed start by

$$H(j_1, \dots, j_\ell) = \langle \{h(j_1, \dots, j_\ell, i_{\ell+1}, \dots, i_n) : n \in \mathbb{N}, n \geq \ell, i_{\ell+1}, \dots, i_n \in \{0, 1\}\} \rangle$$

and the subgroups of $H(j_1, \dots, j_\ell)$ of elements having arguments of minimum length k by

$$H_k(j_1, \dots, j_\ell) = \langle \{h(j_1, \dots, j_\ell, i_{\ell+1}, \dots, i_n) : n \in \mathbb{N}, n \geq k \geq \ell, i_{\ell+1}, \dots, i_n \in \{0, 1\}\} \rangle.$$

Remark 4.2. Let i, j be two sequences such that $h(i)$ and $h(j)$ commute. For any k, ℓ greater than the respective lengths of i, j , we have

$$\langle H_k(i) \cup H_\ell(j) \rangle = H_k(i)H_\ell(j) = H_\ell(j)H_k(i).$$

Let $k \geq 1$ and note that $g_0 h(1)g_0 \notin H$. By using the defining relations for Γ , we compute

$$\begin{aligned} g_0 H_k(0)g_0 &= H_{k+1}(1, 0) \\ h(1)g_0 H_k(0)g_0 h(1) &= h(1)H_{k+1}(1, 0)h(1) = H_{k+1}(1, 1) \\ H \cap g_0 H(1)g_0 &= H(0)H(1, 1) \\ H \cap h(1)g_0 H(1)g_0 h(1) &= h(1)(H \cap g_0 H(1)g_0)h(1) = h(1)H(0)H(1, 1)h(1) = H(0)H(1, 0), \end{aligned}$$

and then

$$\begin{aligned} H \cap g_0 H g_0 &= H(0) H_2(1) \\ H \cap h(1) g_0 H g_0 h(1) &= h(1)(H \cap g_0 H g_0) h(1) = h(1) H(0) H_2(1) h(1) = H(0) H_2(1). \end{aligned}$$

Finally,

$$H \cap g_0 H_k(0) H(1) g_0 = g_0 H_k(0) g_0 (H \cap g_0 H(1) g_0) = H_{k+1}(1, 0) H(0) H(1, 1),$$

where the first equality follows from Dedekind's modular law for groups, and

$$\begin{aligned} H \cap h(1) g_0 H_k(0) H(1) g_0 h(1) &= h(1)(H \cap g_0 H_k(0) H(1) g_0) h(1) \\ &= h(1) H_{k+1}(1, 0) H(0) H(1, 1) h(1) = H_{k+1}(1, 1) H(0) H(1, 0). \end{aligned}$$

Similarly,

$$\begin{aligned} H \cap g_1 H g_1 &= H \cap h(0) g_1 H g_1 h(0) = H(1) H_2(0) \\ H \cap g_1 H(0) H_k(1) g_1 &= H_{k+1}(0, 1) H(1) H(0, 0) \\ H \cap h(0) g_1 H(0) H_k(1) g_1 h(0) &= H_{k+1}(0, 0) H(1) H(0, 1). \end{aligned}$$

Lemma 4.3. *We have $K_0 = H(0)$, $K_1 = H(1)$, and $\ker \Gamma = K_0 \cap K_1 = H(0) \cap H(1) = \{e\}$.*

Proof. We use the notation for $C_{j,k}$ from (2), so that $C_{0,0} = C_{1,0} = H$, and remark that for every j, k we have

$$C_{0,k+1} = \bigcap_{g \in G_0 \setminus H} g C_{1,k} g^{-1} \quad \text{and} \quad C_{1,k+1} = \bigcap_{g \in G_1 \setminus H} g C_{0,k} g^{-1}.$$

In the following we will make use of Remark 4.2, the coset representatives from (7), and the fact that every $C_{j,k}$ is invariant under conjugation by elements of H . We compute that

$$H \cap C_{0,1} = H \cap \bigcap_{g \in G_0 \setminus H} g H g^{-1} = H \cap g_0 H g_0 \cap h(1) g_0 H g_0 h(1) = H(0) H_2(1)$$

and

$$H \cap C_{1,1} = H \cap \bigcap_{g \in G_1 \setminus H} g H g^{-1} = H \cap g_1 H g_1 \cap h(0) g_1 H g_1 h(0) = H(1) H_2(0).$$

Next, let $k \in \mathbb{N}$ and assume $\bigcap_{i=0}^k C_{0,i} = H(0) H_{k+1}(1)$ and $\bigcap_{i=0}^k C_{1,i} = H(1) H_{k+1}(0)$. Then

$$\begin{aligned} \bigcap_{i=0}^{k+1} C_{0,i} &= H \cap \bigcap_{i=0}^k \bigcap_{g \in G_0 \setminus H} g C_{1,i} g^{-1} \\ &= H \cap g_0 \left(\bigcap_{i=0}^k C_{1,i} \right) g_0 \cap h(1) g_0 \left(\bigcap_{i=0}^k C_{1,i} \right) g_0 h(1) \\ &= H \cap g_0 H(1) H_{k+1}(0) g_0 \cap h(1) g_0 H(1) H_{k+1}(0) g_0 h(1) \\ &= H_{k+2}(1, 0) H(0) H(1, 1) \cap H_{k+2}(1, 1) H(0) H(1, 0) \\ &= H(0) H_{k+2}(1), \end{aligned}$$

and similarly

$$\bigcap_{i=0}^{k+1} C_{1,i} = H(1) H_{k+2}(0).$$

Hence, we get that

$$K_0 = \bigcap_{k \geq 0} C_{0,k} = \bigcap_{k \geq 0} H(0) H_{k+1}(1) = H(0)$$

and

$$K_1 = \bigcap_{k \geq 0} C_{1,k} = \bigcap_{k \geq 0} H(1)H_{k+1}(0) = H(1),$$

and thus $\ker \Gamma = K_0 \cap K_1 = H(0) \cap H(1) = \{e\}$. \square

Theorem 4.4. *The group Γ defined above has the unique trace property, but is not C^* -simple.*

Proof. By Lemma 4.3 we have $\ker \Gamma = \{e\}$, so Proposition 3.1 gives that Γ has the unique trace property. Moreover, Γ is countable, H is amenable since it is locally finite, and K_0 and K_1 are nontrivial. Hence, it follows from Proposition 3.5 that Γ is not C^* -simple. \square

Note that Γ contains uncountably many amenable (abelian) subgroups, for example the ones generated by subsets of

$$\{h(\underbrace{0, \dots, 0}_{n \text{ times}}, 1) \mid n \in \mathbb{N}\},$$

so [8, Theorem 1.7] does not apply.

We can find an explicit free nonabelian subgroup of Γ by checking that $\langle g_0 h(1), g_1 h(0) \rangle$ is isomorphic to $\mathbb{Z}_3 * \mathbb{Z}_3$, which clearly contains a nonabelian free group as a subgroup.

Finally, we remark that $\langle K_0 \cup K_1 \rangle = H$, and that the normal closure of H is all of Γ .

Proposition 4.5. *Let*

$$\theta: \Gamma \rightarrow \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$$

be the group homomorphism defined on generators by

$$\theta(h(0)) = (-1, 1), \quad \theta(h(1)) = (1, -1), \quad \theta(g_0) = (1, -1), \quad \theta(g_1) = (-1, 1).$$

Set $\Gamma' = \ker \theta$. Then Γ' is simple.

Proof. Below we only provide the idea of the argument and omit most of the technicalities.

Observe that since $t^{-1}h(0)t = h(0, i_1, \dots, i_{2k})$ for a suitable element $t \in T_{0,2k}$, it follows that $\theta(h(0, i_1, \dots, i_{2k})) = \theta(h(0)) = (-1, 1)$. Therefore

$$h(0)h(0, i_1, \dots, i_{2k}) \in \Gamma'$$

for all $k \in \mathbb{N}$, $j \in \{1, \dots, 2k\}$, and $i_j \in \{0, 1\}$. Analogous arguments yield

$$h(0)h(1, i_1, \dots, i_{2k-1}), h(1)h(1, i_1, \dots, i_{2k}), h(1)h(0, i_1, \dots, i_{2k-1}) \in \Gamma'$$

for all $k \in \mathbb{N}$, $j \in \{1, \dots, 2k\}$, and $i_j \in \{0, 1\}$.

Notice also that $g_1 h(0), h(0)g_1, g_0 h(1), h(1)g_0 \in \Gamma'$. One can now check that

$$\begin{aligned} \Gamma' = \langle & \{h(0)h(0, i_1, \dots, i_{2k}), h(0)h(1, i_1, \dots, i_{2k-1}), h(1)h(1, i_1, \dots, i_{2k}), \\ & h(1)h(0, i_1, \dots, i_{2k-1}), \forall k \in \mathbb{N}, \forall i_j \in \{0, 1\}, \forall j\} \cup \{h(0)g_1, h(1)g_0\} \rangle. \end{aligned}$$

Let $N \neq \{e\}$ be a normal subgroup of Γ' and pick an element $a \in N \setminus \{e\}$. The remainder of the proof is about showing that each of the generators of Γ' listed above can be described by a suitable product of conjugates of a by elements of Γ' . Since these computations are rather tedious, we leave them out. \square

Remark 4.6. If G is an exact group with stable rank one and the unique trace property, then G is C^* -simple by [2, Theorem 2.1]. Let $\Gamma = G_0 *_H G_1$ be the group defined above. Since H is amenable and has finite index in G_0 and G_1 , both these groups are amenable, and therefore Γ is exact by [11, Corollary 3.3]. Hence, Γ does *not* have stable rank one by Theorem 4.4.

Remark 4.7. It was pointed out to us by Adrien Le Boudec [6] that the group Γ is isomorphic to the group $G(A_3, S_3)^*$, which is one of the examples from [4, Section 5].

Let S_3 and A_3 denote the symmetric and alternating group, respectively, on a three-element set. Consider a 3-regular tree T and color the edges $\{1, 2, 3\}$, so that neighboring edges have different colors. Let $\sigma(g, v) \in S_3$ be the permutation of the three colors induced in the natural way by the element $g \in \text{Aut}(T)$. Then $G(A_3, S_3) < \text{Aut}(T)$ is the group of all automorphisms g of T such that $\sigma(g, v) \in A_3$ for all but finitely many vertices v . Then $G(A_3, S_3)^*$ is the subgroup of $G(A_3, S_3)$ of index two preserving the natural bipartition of vertices of T .

To see this, remark that by Bass-Serre's theory (cf. [28, I.4 Theorem 6]) this index two subgroup is an amalgamated product $\bar{G}_0 *_{\bar{H}} \bar{G}_1$, where \bar{G}_0 and \bar{G}_1 are the stabilizers of two adjacent vertices (we will call them v_0 and v_1), and \bar{H} is the stabilizer of the edge between them (we will call it e). Then, using the notation from [5, Subsection 3.1], we may write \bar{G}_i as an increasing union

$$G(S_3)_{v_i} = \bigcup_{n \geq 1}^{\nearrow} K_n(v_i), \quad i = 0, 1,$$

where $K_n(v_i)$ is the subgroup of $G(A_3, S_3)$ that fixes the vertex v_i and has $\sigma(g, v) \in A_3$ for all $g \in K_n(v_i)$ and all v that are at a distance larger than n from v_i . With this notation, the element $h(0, i_2, \dots, i_n)$ acts on T by swapping the two half-trees, emanating from a vertex (we will call it $v(0, i_2, \dots, i_n)$) that is at a distance $n - 1$ from v_0 and at a distance n from v_1 , and not intersecting the geodesic between v_0 and $v(0, i_2, \dots, i_n)$. Clearly, $\sigma(h(0, i_2, \dots, i_n), v(0, i_2, \dots, i_n)) \notin A_3$, because it leaves one edge (therefore one color) fixed. The matching of the other vertices of the half-trees is defined so the local permutations belong to A_3 . This matching is just a matter of orientation of the tree. Adding the element g_1 to the picture, we see that $K_n(v_0)$ is isomorphic to the wreath product $\mathbb{Z}_2 \wr \dots \wr \mathbb{Z}_2 \wr S_3$ ($n - 1$ factors of \mathbb{Z}_2), where the top elements beneath the S_3 factor are $h(1)$, $h(0, 0)$, and $h(0, 1)$. Likewise $K_n(v_1)$ is isomorphic to $\mathbb{Z}_2 \wr \dots \wr \mathbb{Z}_2 \wr S_3$ ($n - 1$ factors of \mathbb{Z}_2), where the top elements beneath the S_3 factor are $h(0)$, $h(1, 0)$ and $h(1, 1)$. In this way, we see that \bar{G}_i is isomorphic to G_i , $i = 0, 1$, and therefore Γ is isomorphic to $G(A_3, S_3)^*$.

The simplicity of the group Γ' is not covered by [5, Corollary 20], because A_3 is not generated by its point stabilizers.

We finally note that $G(A_3, S_3)$ is isomorphic to $\Gamma \rtimes \mathbb{Z}_2$, where \mathbb{Z}_2 acts on Γ by interchanging the indices 0 and 1 of all generating elements of Γ .

5. ACTIONS OF FREE PRODUCTS WITH AMALGAMATION

Let G be any group acting on a space X , and let us first recall some notation. The stabilizer subgroup of an element $x \in X$ is $G_x = \{g \in G : gx = x\}$, and the fixed-point set of $g \in G$ is $X^g = \{x \in X : gx = x\}$. The *kernel* of the action is the set of elements in G acting trivially on X , namely

$$\ker(G \curvearrowright X) = \bigcap_{x \in X} G_x = \{g \in G : X^g = X\}.$$

Note that for every $x \in X$ and all $s, g \in G$, we have $sG_x s^{-1} = G_{sx}$ and $X^{sgs^{-1}} = sX^g$, so the kernel is a normal subgroup of G . The action is called *faithful* when the kernel is trivial, i.e., if for every $g \in G \setminus \{e\}$ there exists $x \in X$ such that $gx \neq x$, and *strongly faithful* if for every finite set $F \subseteq G \setminus \{e\}$ there exists $x \in X$ such that $gx \neq x$ for all $g \in F$.

Furthermore, the action of G on X is called *free* if whenever $g \in G$, $x \in X$, and $gx = x$, then $g = e$. Since $\langle G_x : x \in X \rangle$ is invariant under conjugation, it is a normal subgroup of G , which coincides with the subgroup of G generated by $\{g \in G : X^g \neq \emptyset\}$. This subgroup is the so-called “join” of $\{G_x : x \in X\}$, while $\ker(G \curvearrowright X)$ is the “meet” of $\{G_x : x \in X\}$. Obviously, G acts freely on X if and only if this subgroup is trivial.

Let X be a topological space, and suppose that G acts continuously on X , that is, by homeomorphisms. For every $x \in X$ define G_x^o as the subgroup of G_x consisting of all elements that fix a neighborhood of x pointwise. We notice that $g \in G_x^o$ for some $x \in X$ if and only if X^g has nonempty interior, and we define the *interior* of the action as

$$(8) \quad \text{int}(G \curvearrowright X) = \langle G_x^o : x \in X \rangle = \langle \{g \in G : X^g \text{ has nonempty interior}\} \rangle.$$

Then, by using the identity $X^{sgs^{-1}} = sX^g$, we see that $\{g \in G : X^g \text{ has nonempty interior}\}$ is invariant under conjugation. Indeed, if X^g contains a nonempty open subset V , then sV is a nonempty open subset of $sX^g = X^{sgs^{-1}}$. Therefore, $\text{int}(G \curvearrowright X)$ is automatically a normal subgroup of G . One may also check that $sG_x^o s^{-1} = G_{sx}^o$ for every $x \in X$ and $s \in G$. We say that the action is *topologically free* if X^g has empty interior for every $g \in G \setminus \{e\}$, that is, if the interior is trivial. Finally, we note that for a topological space X , the interior and kernel of an action corresponds to the join and meet, respectively, of $\{G_x^o : x \in X\}$.

Now, fix a nondegenerate amalgam $G = G_0 *_H G_1$ and let T denote its Bass-Serre tree (cf. [28], see also [19] and references therein). Then T has vertex set $G/G_0 \sqcup G/G_1$ and (geometric) edge set G/H . Two vertices in T are adjacent if either of the form

$$(g_0)g_1 \cdots g_{2n-1}G_0 \xrightarrow{(g_0)g_1 \cdots g_{2n-1}g_{2n}H} (g_0)g_1 \cdots g_{2n-1}g_{2n}G_1$$

or of the form (for a transversal edge)

$$(g_0)g_1 \cdots g_{2n}G_1 \xrightarrow{(g_0)g_1 \cdots g_{2n}g_{2n+1}H} (g_0)g_1 \cdots g_{2n}g_{2n+1}G_0$$

for $g_i \in G_{i \pmod{2}} \setminus H$.

We write $d(v, w)$ for the combinatorial distance between two vertices v and w , so that the distance between two adjacent vertices is 1. A *ray* in T is a sequence $(x_n)_{n=0}^\infty$ of vertices, which is geodesic in the sense that $d(x_m, x_n) = |m - n|$ for all m, n (i.e., $x_{n+2} \neq x_n$ for all n). Moreover, given two rays $(x_n)_{n=0}^\infty$ and $(y_n)_{n=0}^\infty$ in T , we say they are *cofinal*, and write $(x_n)_{n=0}^\infty \sim (y_n)_{n=0}^\infty$ if there exist integers k and N such that $y_n = x_{n+k}$ for all $n > N$. Define the boundary ∂T of the Bass-Serre tree T as the set of equivalence classes of cofinal rays.

Henceforth, we fix the sets S_0 and S_1 of representatives for the nontrivial left cosets of G/G_0 and G/G_1 , respectively. Then every $x \in \partial T$ can be uniquely represented by an infinite word $(g_0)g_1g_2 \cdots$, where $g_i \in S_{i \pmod{2}}$, that is, we take either $x_0 = g_0G_1$, $x_1 = g_0g_1G_0$ etc. or $x_0 = G_1$, $x_1 = g_1G_0$, etc., i.e., every class of cofinal rays will be represented by the unique ray in the class starting with gG_1 for $g \in S_0 \cup \{e\}$.

Moreover, ∂T becomes a totally disconnected compact Hausdorff space when equipped with the “shadow topology”, generated by basic open sets $U((g_0)g_1 \cdots g_n)$, where $n \geq 0$ and $g_i \in S_{i \pmod{2}}$, consisting of all equivalence classes of cofinal rays that are identified with infinite words starting with $(g_0)g_1 \cdots g_n$.

The amalgam G acts on its Bass-Serre tree T by left translation, that is, the action of $s \in G$ on vertices is given by $s \cdot gG_0 = sgG_0$, $s \cdot gG_1 = sgG_1$ and on edges by $s \cdot gH = sgH$. Clearly, this also induces an action of G on the boundary of its Bass-Serre tree ∂T .

Recall (3) and define the set

$$(9) \quad K((g_0)g_1 \cdots g_n) = (g_0)g_1 \cdots g_n K_{n+1 \pmod{2}} g_n^{-1} \cdots g_1^{-1} (g_0^{-1}),$$

where $g_i \in S_{i \pmod{2}}$. This is the subgroup of G consisting of all elements that fix the basic open set $U((g_0)g_1 \cdots g_n)$ pointwise.

Lemma 5.1. *We have that $\ker G = \ker(G \curvearrowright T) = \ker(G \curvearrowright \partial T)$, and that $\text{int}(G \curvearrowright \partial T)$ coincides with the normal closure of $K_0 \cup K_1$.*

Proof. It should be clear that $\ker G = \ker(G \curvearrowright T) \subseteq \ker(G \curvearrowright \partial T) \subseteq H$. Suppose that $h \in H \setminus \ker G$. Then there exists g such that $ghg^{-1} \notin H$. Moreover, we can find a g of the

form $(g_0)g_1 \cdots g_n$ with $g_i \in S_{i \pmod{2}}$ with this property. Let x be any ray starting with the corresponding vertices. Then $hx \neq x$, so $h \notin \ker(G \curvearrowright \partial T) \subseteq H$.

Next, K_0 fixes $U(g_0)$, i.e., all sequences of vertices starting with $g_0 G_1$ for any $g_0 \in S_0$, pointwise, and K_1 fixes $U(g_1)$, i.e., all sequences of vertices starting with $G_1, g_1 G_0$ for any $g_1 \in S_1$, pointwise. Hence, $K_0 \cup K_1 \subseteq \text{int}(G \curvearrowright \partial T)$. Therefore, as the latter is normal, the same inclusion holds for the normal closure of $K_0 \cup K_1$.

Suppose that $h \in \text{int}(G \curvearrowright \partial T)$, i.e., $(\partial T)^h$ has nonempty interior. Then h must fix some basic open set pointwise, say $U(g)$ for $g = (g_0)g_1 \cdots g_n$. This means that $h \in K(g)$, as defined in (9), so $h \in gK_{n+1 \pmod{2}}g^{-1}$, that is, h belongs to the normal closure of $K_{n+1 \pmod{2}}$. Since $\text{int}(G \curvearrowright \partial T)$ is generated by $\{h \in G : (\partial T)^h \text{ has nonempty interior}\}$, the conclusion follows.

Note that h also fixes $U((g_0)g_1 \cdots g_{n+1})$, so h belongs to the normal closure of the other K_i as well. In fact, the normal closures of K_0 , K_1 , and $K_0 \cup K_1$ are all the same. \square

Definition 5.2. The subgroup $\text{int}(G \curvearrowright \partial T) = \langle \{g \in G : (\partial T)^g \text{ has nonempty interior}\} \rangle$ of G , or equivalently, the normal closure of $K_0 \cup K_1$ in G , will be called the *interior* of G and denoted $\text{int } G$.

In Proposition 5.4 below, we show that G is C^* -simple if and only if $\text{int } G$ is C^* -simple, giving an analog of Proposition 3.1 (however $\text{int } G$ can be all of G).

Proposition 5.3. *The following are equivalent:*

- (i) $\text{int } G = \{e\}$,
- (ii) $G \curvearrowright T$ is strongly faithful,
- (iii) $G \curvearrowright \partial T$ is strongly faithful,
- (iv) $G \curvearrowright \partial T$ is topologically free, i.e., $G \curvearrowright T$ is slender in the sense of [19].

Proof. The equivalences between (i), (iv), and condition (i) of Theorem 3.2 follow directly from Lemma 5.1 and Definition 5.2. Moreover, condition (ii) of Theorem 3.2 coincides with condition (SF) from [16, p. 245]. Remark that in [16], the notations $\text{Edg}_+ X$ and Y are used for T and ∂T , respectively. In particular, (ii) is the same as condition (SF), and thus the above means that (ii) is equivalent with (i) and (iv). Finally, it follows from [16, Lemma 9] that (ii) implies (iii), and it is stated immediately after the proof of [16, Lemma 9] that its converse holds as well. \square

Using the terminology of [4], we see from the proof of Lemma 5.1 that K_0 and K_1 are precisely the fixators of the “half-trees” of T obtained by removing the edge H .

Proposition 5.4. *Let G be a nondegenerate free product with amalgamation. Then G is C^* -simple if and only if $\text{int } G$ is C^* -simple.*

Proof. Suppose that $\text{int } G$ is C^* -simple.

First, assume that $\text{int } G \subseteq H$. For the moment, write $K_i(G)$ and $K_i(G/\ker G)$ for the K_i ’s corresponding to the amalgams G and $G/\ker G$, respectively. Clearly, a word in G starts with a letter in $G_i \setminus H$ if and only if its image in $G/\ker G$ starts with a letter in $(G_i/\ker G) \setminus (H/\ker G)$. Thus, we see that $K_i(G)/\ker G \cong K_i(G/\ker G)$.

Because $\text{int } G$ is a normal subgroup of G contained in H , we must have $\text{int } G = \ker G$, and then also $K_0 = K_1 = \ker G$. Thus, $G/\text{int } G$ is C^* -simple by Theorem 3.2. Since we have assumed that $\text{int } G$ is C^* -simple, it follows from [8, Theorem 1.4] that G is C^* -simple.

Next, let $C_G(\text{int } G)$ denote the centralizer of $\text{int } G$ in G , and suppose that $g \in C_G(\text{int } G) \setminus H$. In particular, this means that g commutes with all elements in K_0 and K_1 , so $gK_i g^{-1} = K_i$ for $i = 0, 1$. Moreover, for all $g_i \in G_i \setminus H$, we always have that $g_i K_{i+1 \pmod{2}} g_i^{-1} \supseteq K_i$. From this it follows that $K_0 = K_1$, which means that $\ker G = \text{int } G$.

Indeed, let g have length n and denote by g_n the last letter of g . If n is odd, then

$$K_{n \pmod{2}} = gK_{n \pmod{2}}g^{-1} \subseteq \cdots \subseteq g_n K_{n \pmod{2}} g_n^{-1} \subseteq K_{n+1 \pmod{2}},$$

which easily implies $K_0 = K_1$, and hence $\ker G = \text{int } G$. If n is even, then

$$K_{n+1 \pmod 2} = gK_{n+1 \pmod 2}g^{-1} \subseteq \cdots \subseteq g_n K_{n \pmod 2} g_n^{-1} \subseteq K_{n+1 \pmod 2},$$

and therefore all containments are equalities. Assume that $[G_j : H] \geq 3$, for $j = 0$ or 1 , pick a letter $g_j \in G_j \setminus H$ of g , and choose another element $g'_j \in G_j \setminus H$ such that $g_j^{-1}g'_j \in G_j \setminus H$. Then

$$(g'_j)^{-1}g_j K_{j \pmod 2} g_j^{-1}g'_j = (g'_j)^{-1}K_{j+1 \pmod 2}g'_j \supseteq K_{j \pmod 2}.$$

Moreover,

$$(g'_j)^{-1}g_j K_{j \pmod 2} g_j^{-1}g'_j \subseteq K_{j+1 \pmod 2}, \quad \text{so} \quad K_{j \pmod 2} \subseteq K_{j+1 \pmod 2},$$

meaning that $\ker G = \text{int } G$.

Finally, if $C_G(\text{int } G) \subseteq H$, then $C_G(\text{int } G) \subseteq \ker G$, since it is normal in G and contained in H . Then $C_G(\text{int } G) = Z(\text{int } G)$, since it is contained in $\text{int } G$, so it must be trivial since $\text{int } G$ is assumed to be C^* -simple. Hence, it follows from [8, Theorem 1.4] that G is C^* -simple.

The converse holds by [8, Theorem 1.4] because $\text{int } G$ is a normal subgroup of G . \square

Proposition 5.5. *Suppose that $\ker G$ is trivial. Then G is a weak* Powers group.*

Proof. The action of G on the boundary of its Bass-Serre tree is always minimal and strongly hyperbolic (see [19, Proposition 19]). If $\ker G$ is trivial, then the action is also faithful by Lemma 5.1. In [16, Lemma 4] we now replace “strongly faithful” by “faithful”, and the first part of the proof still works, under the assumption that $F \neq \{e\}$ is a one-element set. The rest of the argument goes along the same lines. \square

Example 5.6. The group Γ from Section 4 is a weak* Powers group that is not C^* -simple.

We complete this section by some facts about boundary actions and refer to [8, 21, 25] for further details. An action of G on a compact space X is called a *boundary action* if it is minimal (i.e., the orbits are dense) and strongly proximal (i.e., the orbit-closure in $P(X)$ of every probability measure on X contains a point mass). In this case we also say that the space X is a G -boundary.

For every group G there is a universal G -boundary called the *Furstenberg boundary* and denoted $\partial_F G$. If X is any other G -boundary, then there exists a continuous surjective G -equivariant map $\partial_F G \rightarrow X$. Moreover, we remark that for every $g \in G$, the set $(\partial_F G)^g$ is always clopen, cf. [8, Lemma 3.3].

In Section 7 we will often make use of the following observation from [8].

Lemma 5.7. *Let G be a group, N a normal subgroup of G , and L a normal subgroup of N . Suppose that $g \in N$ is such that $(\partial_F G)^g \neq \emptyset$. Then $(\partial_F(N/L))^{g^L} \neq \emptyset$.*

Proof. First, the quotient map $N \rightarrow N/L$ gives rise to an action of N on $\partial_F(N/L)$, which makes $\partial_F(N/L)$ an N -boundary. Thus, there exists an N -equivariant continuous surjective map $\psi: \partial_F N \rightarrow \partial_F(N/L)$. Next, [8, Lemma 5.2] says that the N -action on $\partial_F N$ extends to an action of G , such that $\partial_F N$ becomes a G -boundary. Therefore, there is a G -equivariant continuous surjective map $\phi: \partial_F G \rightarrow \partial_F N$, so altogether we have surjections

$$\partial_F G \xrightarrow{\phi} \partial_F N \xrightarrow{\psi} \partial_F(N/L).$$

Now, if $g \in N$ and $(\partial_F G)^g \neq \emptyset$, there exists $x \in \partial_F G$ such that $gx = x$. It follows that $g\psi(\phi(x)) = \psi(\phi(gx)) = \psi(\phi(x))$, so $\psi(\phi(x)) \in (\partial_F(N/L))^g = (\partial_F(N/L))^{g^L}$, which is therefore nonempty. In particular, we have $\psi(\phi((\partial_F G)^g)) \subseteq (\partial_F(N/L))^{g^L}$. \square

Lemma 5.8. *Suppose that G is a nondegenerate amalgam. Then ∂T is a G -boundary.*

Proof. This follows from [25, Example 2] and [19, Proof of Proposition 13]. \square

Theorem 5.9. *Suppose that K_0 or K_1 is amenable. Then G is C^* -simple if and only if both K_0 and K_1 are trivial.*

Proof. Assume first that G is C^* -simple. Let $x \in \partial T$ be identified with a sequence of vertices $g_0 G_1, g_0 g_1 G_0$, etc., where $g_i \in S_{i \pmod{2}}$ (of course, the argument is similar if it starts with $G_1, g_1 G_0$, etc). Then G_x^o is the direct limit of the sequence $K(g_0) \subseteq K(g_0 g_1) \subseteq \cdots$, that is, of

$$K_0 \subseteq g_0 K_1 g_0^{-1} \subseteq g_0 g_1 K_0 g_1^{-1} g_0^{-1} \subseteq \cdots$$

Clearly, K_0 is amenable if and only if K_1 is amenable, since either of them is a subgroup of a conjugate of the other, see (5). Therefore, all $K((g_0)g_1 \cdots g_n)$ are also amenable, since they are conjugates of K_0 or K_1 , see (9). As the class of amenable groups is closed under direct limits, we have that G_x^o is amenable.

Since ∂T is a G -boundary by Lemma 5.8 and G is assumed to be C^* -simple, [8, Corollary 7.5] says that $C(\partial T) \rtimes_r G$ is simple. Then it follows from [25, Proposition 14 (ii)] that $G \curvearrowright \partial T$ is topologically free. Hence, Proposition 5.3 gives that $\text{int } G = \{e\}$, that is, both K_0 and K_1 are trivial.

Conversely, if $K_0 = K_1 = \{e\}$, then G is C^* -simple by Theorem 3.2. \square

A similar argument shows that G_x^o is C^* -simple for all $x \in \partial T$ if K_0 or K_1 is C^* -simple, and it seems likely that this implies C^* -simplicity of G .

Remark 5.10. Any amalgamated free product where K_0 and K_1 are nontrivial, amenable, and $K_0 \cap K_1 = \{e\}$ is not C^* -simple, but has the unique trace property. As noted in Remark 4.7, the group Γ of Section 4 is isomorphic to one of Le Boudec's examples from [4, Section 5]. However, if the groups G_0 and G_1 are nonisomorphic, then such an amalgamated product will not be covered by [4, Section 5].

6. RADICAL AND RESIDUAL CLASSES OF GROUPS

In this section, by a *class of groups*, we will always mean a class X of groups that contains the trivial group and is closed under isomorphisms, i.e., if $G \in X$ and $H \cong G$, then $H \in X$. In [27], this is called a *group theoretical class*.

Let X be a class of groups and let G be any group. Define $\rho(G)$ as the normal subgroup of G generated by all normal subgroups of G that belong to X , and $\rho^*(G)$ as the intersection of all normal subgroups of G with quotient belonging to X , i.e.,

$$\rho(G) = \prod \{N : N \triangleleft G \text{ and } N \in X\},$$

$$\rho^*(G) = \bigcap \{N : N \triangleleft G \text{ and } G/N \in X\}.$$

These are both normal subgroups of G , called join and meet of the respective families. Whenever more than one class is around, we will often write ρ_X and ρ_X^* .

Definition 6.1. A class of groups X is called a *radical* class if it is closed under quotients (i.e., closed under homomorphic images), and if for any group G we have

- (i) $\rho(G) \in X$,
- (ii) $\rho(G/\rho(G)) = \{e\}$.

A class of groups X is called a *residual* (or *coradical* or *semisimple*) class if it is closed under normal subgroups, and if for any group G we have

- (i*) $G/\rho^*(G) \in X$,
- (ii*) $\rho^*(\rho^*(G)) = \rho^*(G)$.

Clearly, if X is radical, then for any G we have $\rho(\rho(G)) = \rho(G)$, and $G \in X$ if and only if $\rho(G) = G$. Moreover, if X is residual, then for any G we have $\rho^*(G/\rho^*(G)) = \{e\}$, and $G \in X$ if and only if $\rho^*(G) = \{e\}$.

The above definitions are not completely consistent within references; some say that a class is radical if it is closed under quotients and (i) holds, and *strict* radical if (ii) holds as well (similarly for residual), and there are possibly other variations.

Proposition 6.2. *A class of groups is radical if and only if it is closed under quotients, extensions, and satisfies (i). Moreover, a class of groups is residual if and only if it is closed under normal subgroups, extensions, and satisfies (i*).*

Proof. This is explained in [27, Theorems 1.32 and 1.35], where (i) and (i*) are equivalent to the properties $NX = X$ and $RX = X$ of [27, p. 20 and p. 23], respectively, see [27, p. 19]. \square

Let X be a class of groups closed under quotients, and define X^* to be all groups satisfying $\rho(G) = \{e\}$, that is, the class of groups with no nontrivial normal subgroups in X , i.e.,

$$X^* = \{G : \rho(G) = \{e\}\} = \{G : N \triangleleft G, N \neq \{e\} \Rightarrow N \notin X\}.$$

Similarly, if X is a class of groups closed under normal subgroups, define X_* to be all groups satisfying $\rho^*(G) = G$, that is, the class of groups with no nontrivial quotients in X , i.e.,

$$X_* = \{G : \rho^*(G) = G\} = \{G : N \triangleleft G, N \neq G \Rightarrow G/N \notin X\}.$$

Proposition 6.3. *Let X be a class of groups closed under quotients. Then X is radical if and only if $X = (X^*)_*$ if and only if $X = Y_*$ for some class of groups Y that is closed under normal subgroups.*

Let X be a class of groups closed under normal subgroups. Then X is residual if and only if $X = (X_)^*$ if and only if $X = Y^*$ for some class of groups Y that is closed under quotients.*

Proof. This follows from [27, Theorems 1.38 and 1.39], see [27, p. 6-7] for notation. \square

Example 6.4. Let X be the class of amenable groups, which is known to be a radical class. Then [8, Theorem 1.3] shows that the class of groups with the unique trace property coincides with X^* , and is therefore residual by Proposition 6.3. Thus, a group is amenable if and only if it does not have any nontrivial quotient with the unique trace property.

Example 6.5. Let X be the class of Powers (resp. weak Powers, weak* Powers) groups, which is not closed under extensions, so it is not a residual class. However, X is closed under normal subgroups by Lemma 2.2, meaning that we can define the class X_* of groups with no nontrivial quotient which is a Powers (resp. weak Powers, weak* Powers) group. Moreover, X_* is a radical class, but $\rho_{X_*}(G) = \{e\}$ does *not* imply that $G \in X$.

If X is the class of all Powers groups, then $P = (X_*)^*$ is the residual closure of X , that is, the smallest residual class containing all Powers groups. In Section 7 we will see that the class of C^* -simple groups is residual, and hence it contains P (it could possibly coincide with P).

Example 6.6. Some radical classes of groups are e.g. locally finite groups, elementary amenable groups, amenable groups, and groups that do not contain any nonabelian free subgroup. We denote the latter class by NF , which gives rise to the residual class $AF = (NF)^*$, consisting of all groups for which every nontrivial normal subgroup contains a free nonabelian subgroup, and $G \in NF$ if and only if $\rho_{AF}^*(G) = G$. Since every amenable group belongs to NF , every group in AF has the unique trace property. Moreover, (1) gives that every amalgamated free product with trivial kernel belongs to AF .

Both the class of amenable groups and NF are closed under subgroups, but in general, radical classes are not necessarily closed even under normal subgroups. A radical class that is closed under normal subgroups is sometimes called *hereditary*.

Lemma 6.7. *Let X be a class of groups satisfying (i). Then X is closed under normal subgroups if and only if for any group G and normal subgroup N of G we have*

$$\rho(N) = \rho(G) \cap N.$$

In particular, this implies that $\rho(N)$ is a normal subgroup of G .

Proof. If $G \in X$ and N is normal in G , then $\rho(N) = \rho(G) \cap N = G \cap N = N$, so $N \in X$.

The converse is explained in [27, Lemma 1.31, Corollaries 1 and 2]. \square

A direct proof in the special case of countable amenable groups is given in [29, Corollary B.4]. The result below is similar to [29, Corollary B.6], using Lemma 6.7.

For a group G and a subset N , we let $Z_G(N)$ and $Z(G)$ denote the centralizer of N in G , and the center of G , respectively.

Lemma 6.8. *Let X be a class of groups that satisfies (i) and is closed under normal subgroups. Assume that G is any group and N is a normal subgroup of G such that $\rho(N) = \{e\}$. Then $\rho(G) = \rho(Z_G(N))$.*

Proof. Since $\{e\} = \rho(N) = \rho(G) \cap N$, the normal subgroups $\rho(G)$ and N commute, so $\rho(G) \subseteq Z_G(N)$. Hence, $\rho(Z_G(N)) = \rho(G) \cap Z_G(N) = \rho(G)$. \square

Lemma 6.9. *Let X be a class of groups that satisfies (i), is closed under normal subgroups, and contains all abelian groups. Assume that G is any group and N is a normal subgroup of G such that both $\rho(N)$ and $\rho(G/N)$ are trivial. Then $\rho(G) = \{e\}$.*

Proof. First, $Z(N)$ is normal in N , so Lemma 6.7 gives that $\rho(Z(N)) \subseteq \rho(N) = \{e\}$, and since $Z(N) \in X$, we must have $Z(N) = \{e\}$. Thus, the map $Z_G(N) \rightarrow G/N$, $x \mapsto xN$ is injective, and $\rho(G/N) = \{e\}$ implies $\rho(Z_G(N)) = \{e\}$. By Lemma 6.8, we thus get $\rho(G) = \{e\}$. \square

Lemma 6.10. *Let X be a residual class, and suppose that X_* is closed under normal subgroups and contains all abelian groups. Let G be any group and N a normal subgroup of G . Then G belongs to X if and only if both N and $Z_G(N)$ belong to X .*

Proof. Clearly, X_* satisfies the conditions of Lemmas 6.8 and 6.9, and by Proposition 6.3 we know that $G \in X$ if and only if $\rho_{X_*}(G) = \{e\}$. \square

The above result is an analog of [8, Theorem 1.4], while the result below shows that when X is a residual class and G is a group, then $\rho_{X_*}(G)$ is the smallest normal subgroup of G that produces a quotient in X .

Lemma 6.11. *Let X be a class of groups closed under normal subgroups and suppose that G is a group with a normal subgroup N such that $G/N \in X$. Then $\rho_{X_*}(G) \subseteq N$.*

Proof. If $G/N \in X$, then $\rho_{X_*}(G/N) = \{e\}$. Suppose that $\rho_{X_*}(G)$ is not contained in N . Then $\rho_{X_*}(G)/(\rho_{X_*}(G) \cap N)$ is isomorphic to $(\rho_{X_*}(G)N)/N$, which is a normal nontrivial subgroup of G/N that belongs to X_* . This is a contradiction. \square

Lemma 6.12. *Let X be a class of groups that is closed under normal subgroups, extensions, and contains all finite groups. Assume that G is any group and H is a subgroup of G of finite index. Then $G \in X$ if and only if $H \in X$.*

Proof. Let N be the normal core of H in G , that is, the largest normal subgroup of G contained in H . It is well-known that N also has finite index in G . Suppose first that $G \in X$. Then $N \in X$ and H/N is finite, so $H/N \in X$, and hence $H \in X$. The converse is similar; if $H \in X$, then $N \in X$ and G/N is finite, so $G \in X$. \square

7. AMENABLISH GROUPS AND THE AMENABLISH RADICAL

Since the class of C^* -simple groups is closed under normal subgroups [8, Theorem 1.4], the following definition makes sense in light of Proposition 6.3.

Definition 7.1. We call a group *amenablisth* if it has no nontrivial C^* -simple quotients. The class of all amenablisth groups is radical, so every group G has a unique maximal normal amenablisth subgroup, which will be called the *amenablisth radical* of G .

It is clear that every amenable group is amenablisth, but not all amenablisth groups are amenable, as explained in Corollary 7.11 below.

We will now show that the class of C^* -simple groups is residual, which will imply that a group is C^* -simple precisely when its amenablisth radical is trivial. Since the class of C^* -simple groups is known to be closed under normal subgroups and extensions by [8, Theorem 1.4], we only have to prove that (i*) from Definition 6.1 holds.

Proposition 7.2. *Suppose that G is a group and $\{N_\alpha\}_{\alpha \in \Lambda}$ is a family of normal subgroups of G such that G/N_α is C^* -simple for all α .*

Then $G/\bigcap_{\alpha \in \Lambda} N_\alpha$ is C^ -simple.*

Proof. For any two indices α and β , the group $(N_\alpha N_\beta)/N_\beta \cong N_\alpha/(N_\alpha \cap N_\beta)$ is a normal subgroup of G/N_β , so it is C^* -simple again by [8, Theorem 1.4]. Moreover, $G/(N_\alpha \cap N_\beta) \rightarrow G/N_\alpha$ is surjective with kernel $N_\alpha/(N_\alpha \cap N_\beta)$. Hence, applying [8, Theorem 1.4] once more gives that $G/(N_\alpha \cap N_\beta)$ is C^* -simple. We may therefore assume that the family $\{N_\alpha\}_{\alpha \in \Lambda}$ is closed under finite intersections.

It is easy to see, using transfinite induction and the Axiom of Choice, that we can obtain a well-ordered set $\{N_\beta\}_{\beta \in I}$ of normal subgroups of G with the property

$$\bigcap_{\beta \in I} N_\beta = \bigcap_{\alpha \in \Lambda} N_\alpha \stackrel{\text{def}}{=} N.$$

After factoring the whole family by N , we deduce the following equivalent reformulation of the above statement: Suppose G is a group with a decreasing (transfinite) sequence

$$G = N_0 \supsetneq N_1 \supsetneq N_2 \supsetneq \cdots \supsetneq N_\alpha \supsetneq \cdots \supsetneq N_\beta \supsetneq \cdots$$

that satisfies

- (i) N_α is normal in G for all $\alpha \in I$,
- (ii) $\bigcap_{\alpha \in I} N_\alpha = \{e\}$,
- (iii) G/N_α is C^* -simple for all $\alpha \in I$.

Then G is C^* -simple.

To prove the latter statement, set $X_\alpha = \partial_F(N_\alpha/N_{\alpha+1})$, where ∂_F denotes the Furstenberg boundary. Then X_α is a G -boundary for all α . Indeed, by [8, Lemma 5.2] and the normality assumption, the action of $N_\alpha/N_{\alpha+1}$ on X_α extends uniquely to a boundary action of $G/N_{\alpha+1}$ on X_α . Then, composition with the quotient map gives a boundary action of G on X_α (compare with [8, Proof of Theorem 1.4]).

Now, set $X = \prod_{\alpha} X_\alpha$ (with the usual product topology). We wish to show that the action of G on X is a boundary action, that is, strongly proximal and minimal. First, [25, Lemma 3] gives that the action of G on X is strongly proximal, since it is strongly proximal on each factor. Next, take an arbitrary point $x = (x_\alpha) \in X$ and an arbitrary basic open set $U = \prod_{\alpha} U_\alpha$, where we have $U_\alpha = X_\alpha$ except for finitely many sets $U_{\alpha_1}, \dots, U_{\alpha_n}$, with $\alpha_1 < \dots < \alpha_n$. Note that for all α , the set $(N_\alpha \setminus N_{\alpha+1})x_\alpha$ is dense in X_α and also that N_α acts minimally on X_α , while $N_{\alpha+1}$ acts trivially on X_α . Therefore there exists a group element $g_1 \in N_{\alpha_1}$, such that $g_1 x_{\alpha_1} \in U_{\alpha_1}$. Also there exists $g_2 \in N_{\alpha_2}$, such that $g_2 g_1 x_{\alpha_2} \in U_{\alpha_2}$. We continue the argument and finally deduce that there exists $g_n \in N_{\alpha_n}$ with $g_n \cdots g_1 x_{\alpha_n} \in U_{\alpha_n}$. This shows that $g_n \cdots g_1 x \in U$. We conclude that Gx is dense in X , hence, X is a G -boundary. Note that the last argument does not require the Axiom of Choice, since we just need the existence of g_i 's and not their concrete choice.

We wish to show that this action is free, i.e., that for all nontrivial $g \in G$, the set $X^g = \{x : gx = x\}$ is empty. So pick an arbitrary $g \in G \setminus \{e\}$. From the assumption it follows that there must be a unique α such that $g \in N_\alpha \setminus N_{\alpha+1}$. Clearly, $N_\alpha/N_{\alpha+1}$ is C^* -simple, since it is a normal subgroup of $G/N_{\alpha+1}$, which is C^* -simple by assumption, so [21, Theorem 6.2] and [8, Lemma 3.3] imply that $N_\alpha/N_{\alpha+1}$ acts freely on $\partial_F(N_\alpha/N_{\alpha+1})$, that is,

$$X_\alpha^g = \partial_F(N_\alpha/N_{\alpha+1})^{gN_{\alpha+1}} = \emptyset$$

Hence, $X^g = \prod_\alpha X_\alpha^g = \emptyset$.

The result now follows from [21, Theorem 6.2]. \square

In Proposition 7.2 the use of the Axiom of Choice can be avoided for groups that are concretely given, e.g. if G is given by a totally ordered set of generators and relations, or if G is given by some concrete dynamical properties.

Corollary 7.3. *The class of C^* -simple groups is a residual class.*

Hence, a group G is C^ -simple if and only if its amenablsh radical N is trivial, and N is the smallest normal subgroup of G that produces a C^* -simple quotient.*

We will now describe the amenablsh radical in terms of the Furstenberg boundary.

Definition 7.4. Let G be any group. Set $N_0 = \{e\}$ and $N_1 = \text{int}(G \curvearrowright \partial_F G)$, recall (8), and moreover, for every ordinal α , define a normal subgroup $N_{\alpha+1}$ of G by

$$N_{\alpha+1}/N_\alpha = \text{int}(G/N_\alpha \curvearrowright \partial_F(G/N_\alpha)),$$

and for every limit ordinal β , set $N_\beta = \bigcup_{\alpha < \beta} N_\alpha$, which is clearly also normal in G . Then $\{N_\alpha\}_\alpha$ is an ascending normal series of G which eventually stabilizes, and we finally set $AH(G) = \bigcup_\alpha N_\alpha$.

Lemma 7.5. *For any group G , the quotient $G/AH(G)$ is C^* -simple.*

Proof. Let $\{N_\alpha\}_\alpha$ be as in Definition 7.4. Then there exists some ordinal β such that $AH(G) = N_\beta$. If G/N_β is not C^* -simple, then $N_{\beta+1}/N_\beta = (\partial_F(G/N_\beta))^{gN_\beta} \neq \emptyset$ is nontrivial. Hence, $N_\beta \subsetneq N_{\beta+1}$, contradicting the definition of $AH(G)$. \square

Note that $G/\text{int}(G \curvearrowright \partial_F G)$ is not necessarily C^* -simple, i.e., $AH(G)$ is in general bigger than $\text{int}(G \curvearrowright \partial_F G)$. Indeed, it was explained to us [6] that by applying [7, Theorem 1.11], one can construct an example $G = G(F, F')$ such that $\text{int}(G \curvearrowright \partial_F G)$ has index two in G (the condition is that F' is generated by its point stabilizers). We refer to [5, 4, 7] for more about this construction.

Lemma 7.6. *Suppose that N is a normal subgroup of a group G such that G/N is C^* -simple. Then $AH(G) \subseteq N$.*

Proof. The action of G/N on $X = \partial_F(G/N)$ is free by [21, Theorem 6.2]. Pick $g \in \text{int}(G \curvearrowright \partial_F G)$ such that $(\partial_F G)^g \neq \emptyset$. By Lemma 5.7, we have $X^{gN} \neq \emptyset$, which means that gN is trivial in G/N , i.e., $g \in N$. Since the set of all g with $(\partial_F G)^g \neq \emptyset$ generates $\text{int}(G \curvearrowright \partial_F G)$, it follows that $\text{int}(G \curvearrowright \partial_F G) \subseteq N$.

We continue by transfinite induction. Let $\{N_\alpha\}_\alpha$ be the series from Definition 7.4 associated with G . We have shown that $N_1 \subseteq N$. Assume next that $N_\alpha \subseteq N$ for some ordinal α and note that there is a quotient map $G/N_\alpha \rightarrow G/N$. Choose $g \in N_{\alpha+1}$ such that $(\partial_F(G/N_\alpha))^{gN_\alpha} \neq \emptyset$. Then the same argument as above gives that $X^{gN} \neq \emptyset$, so $g \in N$. Hence, we conclude that $N_{\alpha+1} \subseteq N$. Finally, if β is a limit ordinal and $N_\alpha \subseteq N$ for all $\alpha < \beta$, then clearly $N_\beta \subseteq N$. \square

Lemma 7.7. *Let N be a normal subgroup of G . Then $AH(N) = AH(G) \cap N$.*

Proof. Pick $g \in \text{int}(G \curvearrowright \partial_F G) \cap N$ such that $(\partial_F G)^g \neq \emptyset$. It follows from Lemma 5.7 that $(\partial_F N)^g \neq \emptyset$, so $g \in \text{int}(N \curvearrowright \partial_F N)$. Since the set of all g with $(\partial_F G)^g \neq \emptyset$ generates $\text{int}(G \curvearrowright \partial_F G)$, we get that $\text{int}(G \curvearrowright \partial_F G) \cap N \subseteq \text{int}(N \curvearrowright \partial_F N)$.

We continue by transfinite induction. Let $\{N_\alpha\}_\alpha$ and $\{H_\alpha\}_\alpha$ be the series from Definition 7.4 associated with G and N , respectively. We have shown that $N_1 \cap N \subseteq H_1$. Let α be an ordinal number and assume that $N_\alpha \cap N \subseteq H_\alpha$. Note that $N/(N \cap N_\alpha) \cong (NN_\alpha)/N_\alpha$ is a normal subgroup of G/N_α , and that N/H_α is a quotient of $N/(N \cap N_\alpha)$. Choose $g \in N_{\alpha+1} \cap N$ such that $\partial_F(G/N_\alpha)^{gN_\alpha} \neq \emptyset$. Then by Lemma 5.7 we have $(\partial_F(N/H_\alpha))^{gH_\alpha} \neq \emptyset$. Hence, $gH_\alpha \in \text{int}(N/H_\alpha \curvearrowright \partial_F(N/H_\alpha)) = H_{\alpha+1}/H_\alpha$, and it follows that $N_{\alpha+1} \cap N \subseteq H_{\alpha+1}$. Finally, if β is a limit ordinal and $N_\alpha \cap N \subseteq H_\alpha$ for all $\alpha < \beta$, then clearly $N_\beta \cap N \subseteq H_\beta$. Thus, we have $AH(G) \cap N \subseteq AH(N)$.

For the opposite inclusion, note that $G/AH(G)$ is C^* -simple by Lemma 7.5, and that $(AH(G)N)/AH(G)$ is normal in $G/AH(G)$, so $N/(AH(G) \cap N) \cong (AH(G)N)/AH(G)$ is C^* -simple by using [8, Theorem 1.4]. Hence, Lemma 7.6 gives that $AH(N) \subseteq AH(G) \cap N$. \square

Proposition 7.8. *For any group G , the amenablsh radical of G coincides with $AH(G)$.*

Proof. We need to show that $AH(G)$ is amenablsh, and that it contains all normal amenablsh subgroups of G .

Set $M = \text{int}(G \curvearrowright \partial_F G)$. Suppose first that L is a normal subgroup of M such that $L \neq M$. Pick $g \in M \setminus L$ so that $(\partial_F G)^g \neq \emptyset$. It follows from Lemma 5.7 that $(\partial_F(M/L))^{gL} \neq \emptyset$, so M/L is not C^* -simple. Hence, M is amenablsh.

Let $\{N_\alpha\}_\alpha$ be the series from Definition 7.4 associated with G . Then it follows that $N_{\alpha+1}/N_\alpha$ is amenablsh for every ordinal α , by using the same argument as above with G/N_α in place of G . Since the class of amenablsh groups is radical, it is closed under extensions and under increasing unions of normal subgroups. Since N_1 is amenablsh, an argument by transfinite induction gives that N_α is amenablsh for every ordinal α . Hence, $AH(G)$ is amenablsh.

Next, let L be an amenablsh normal subgroup of G , and assume that L is not contained in $AH(G)$. Set $K = L \cap AH(G)$, then $K \neq L$ and $K = AH(L)$ by Lemma 7.7. Hence, L/K is C^* -simple by Lemma 7.5. \square

Lemma 7.9. *The class of amenablsh groups is closed under normal subgroups.*

Proof. This is a direct consequence of Lemma 6.7, Lemma 7.7, and Proposition 7.8. \square

Lemma 7.10. *Let G be any group and H a subgroup of finite index. Then G is amenablsh if and only if H is amenablsh.*

Proof. The class of amenablsh groups is closed under normal subgroups, extensions, and contains all finite groups. Hence, Lemma 6.12 applies. \square

Corollary 7.11. *The group Γ of Section 4 is amenablsh, but not amenable (and has trivial amenable radical).*

Proof. By Theorem 4.4, the group Γ is not C^* -simple, but has the unique trace property, so it has trivial amenable radical and is icc. The normal subgroup Γ' from Proposition 4.5 is not C^* -simple either, because it has finite index in Γ (see [17, Proposition 19 (iv)]). Since Γ' is simple and $AH(\Gamma') \neq \{e\}$, we must have $AH(\Gamma') = \Gamma'$, that is, Γ' is amenablsh. Hence, it follows from Lemma 7.10 that Γ is amenablsh. \square

Remark 7.12. The class of amenablsh groups is not closed under subgroups. Indeed, by Corollary 7.11 the group Γ is amenablsh, but it contains as subgroup a nonabelian free group, which is C^* -simple, and thus not amenablsh.

Moreover, $AH(\Gamma) = \Gamma$ and $NF(\Gamma) = \{e\}$, while a recent preprint by Olshanskii and Osin [24] presents a group G with the property $AH(G) = \{e\}$ and $NF(G) = G$. Hence, it seems to be no relation between the class of amenable groups and NF (except that both contain all amenable groups).

Remark 7.13. Let G be a countable group, and let N be the subgroup of G generated by all recurrent amenable subgroups in G (see [22, Definition 1.4]). Does N coincide with $AH(G)$?

It is mentioned in [22, Remark 6.4] that for every $x \in \partial_F G$ the subgroup G_x is recurrent amenable in G , so $\text{int}(G \curvearrowright \partial_F G) = \langle G_x : x \in \partial_F G \rangle \subseteq N$. Moreover, [22, Theorem 6.3] says that N is trivial if and only if G is C^* -simple, that is, if and only if $\text{int}(G \curvearrowright \partial_F G)$ is trivial.

Note also that a recent preprint [7, Theorem 2.16] introduces a unique maximal uniformly recurrent subgroup \mathcal{A}_G of G , and [7, Proposition 2.20 (ii)] states that $\langle H : H \in \mathcal{A}_G \rangle = \text{int}(G \curvearrowright \partial_F G)$, which in general is smaller than $AH(G)$, cf. comment after Lemma 7.5.

Example 7.14. If G is a simple group, then G is either C^* -simple or amenable. E.g. Thompson's group T is known to be simple, so it follows directly from [7, 15] that T is amenable if and only if Thompson's group F is amenable.

ACKNOWLEDGEMENTS

The second author is funded by the Research Council of Norway through FRINATEK, project no. 240913. Part of the work of the second author was conducted during a research stay at Arizona State University, Fall 2015, and another part when attending the research program "Classification of operator algebras: complexity, rigidity, and dynamics" at the Mittag-Leffler Institute outside Stockholm, January–February 2016. In both cases, he would like to thank the hosts for providing excellent working conditions. The second author is also grateful to Erik Bédos for useful feedback on an early version of the article. Finally, the authors would like to thank the referee for several valuable comments.

REFERENCES

- [1] Erik Bédos. Operator algebras associated with free products of groups with amalgamation. *Math. Ann.*, 266(3):279–286, 1984.
- [2] Erik Bédos and Tron Omland. On reduced twisted group C^* -algebras that are simple and/or have a unique trace. Preprint arXiv:1606.02637, 2016.
- [3] Florin P. Boca and Viorel Nițică. Combinatorial properties of groups and simple C^* -algebras with a unique trace. *J. Operator Theory*, 20(1):183–196, 1988.
- [4] Adrien Le Boudec. C^* -simplicity and the amenable radical. *Invent. Math.*, to appear.
- [5] Adrien Le Boudec. Groups acting on trees with almost prescribed local action. *Comment. Math. Helv.*, 91(2):253–293, 2016.
- [6] Adrien Le Boudec. E-mail correspondence, 2016.
- [7] Adrien Le Boudec and Nicolás Matte Bon. Subgroup dynamics and C^* -simplicity of groups of homeomorphisms. Preprint arXiv:1605.01651, 2016.
- [8] Emmanuel Breuillard, Mehrdad Kalantar, Matthew Kennedy, and Narutaka Ozawa. C^* -simplicity and the unique trace property for discrete groups. Preprint arXiv:1410.2518, 2014.
- [9] Yves de Cornulier. Infinite conjugacy classes in groups acting on trees. *Groups Geom. Dyn.*, 3(2):267–277, 2009.
- [10] Mahlon M. Day. Amenable semigroups. *Illinois J. Math.*, 1:509–544, 1957.
- [11] Kenneth J. Dykema. Exactness of reduced amalgamated free product C^* -algebras. *Forum Math.*, 16(2):161–180, 2004.
- [12] Kenneth J. Dykema, Uffe Haagerup, and Mikael Rørdam. The stable rank of some free product C^* -algebras. *Duke Math. J.*, 91(1):95–121, 1997.
- [13] Kenneth J. Dykema and Pierre de la Harpe. Some groups whose reduced C^* -algebras have stable rank one. *J. Math. Pures Appl. (9)*, 78(6):591–608, 1999.
- [14] Uffe Haagerup. A new look at C^* -simplicity and the unique trace property of a group. In *Operator algebras and applications*, vol. 12 of *Abel Symp.*, pp. 161–170. Springer, Heidelberg, 2016.
- [15] Uffe Haagerup and Kristian Knudsen Olesen. The Thompson groups T and V are not inner amenable. Preprint arXiv:1609.05086, 2016.

- [16] Pierre de la Harpe. Reduced C^* -algebras of discrete groups which are simple with a unique trace. In *Operator algebras and their connections with topology and ergodic theory (Buşteni, 1983)*, vol. 1132 of *Lecture Notes in Math.*, pp. 230–253. Springer, Berlin, 1985.
- [17] Pierre de la Harpe. On simplicity of reduced C^* -algebras of groups. *Bull. Lond. Math. Soc.*, 39(1):1–26, 2007.
- [18] Pierre de la Harpe and K. Jhabvala. Quelques propriétés des algèbres d'un groupe discontinu d'isométries hyperboliques. In *Ergodic theory (Sem., Les Plans-sur-Bex, 1980)*, vol. 29 of *Monograph. Enseign. Math.*, pp. 47–55. Univ. Genève, Geneva, 1981.
- [19] Pierre de la Harpe and Jean-Philippe Préaux. C^* -simple groups: amalgamated free products, HNN extensions, and fundamental groups of 3-manifolds. *J. Topol. Anal.*, 3(4):451–489, 2011.
- [20] Nikolay A. Ivanov. On the structure of some reduced amalgamated free product C^* -algebras. *Internat. J. Math.*, 22(2):281–306, 2011.
- [21] Mehrdad Kalantar and Matthew Kennedy. Boundaries of reduced C^* -algebras of discrete groups. *J. Reine Angew. Math.*, to appear.
- [22] Matthew Kennedy. Characterizations of C^* -simplicity. Preprint arXiv:1509.01870, 2015.
- [23] Sang Og Kim. Powers groups and crossed product C^* -algebras. *Commun. Korean Math. Soc.*, 5(2):147–149, 1990.
- [24] Alexander Yu. Olshanskii and Denis V. Osin. C^* -simple groups without free subgroups. *Groups Geom. Dyn.*, 8(3):933–983, 2014.
- [25] Narutaka Ozawa. Lecture on the Furstenberg boundary and C^* -simplicity, 2014. <http://www.kurims.kyoto-u.ac.jp/~narutaka/notes/yokou2014.pdf>.
- [26] Robert T. Powers. Simplicity of the C^* -algebra associated with the free group on two generators. *Duke Math. J.*, 42:151–156, 1975.
- [27] Derek J. S. Robinson. *Finiteness conditions and generalized soluble groups. Part 1*, volume 62 of *Ergeb. Math. Grenzgeb. (2)*. Springer-Verlag, New York-Berlin, 1972.
- [28] Jean-Pierre Serre. *Arbres, amalgames, SL_2* , volume 46 of *Astérisque*. Soc. Math. France, Paris., 1977.
- [29] Robin D. Tucker-Drob. Shift-minimal groups, fixed price 1, and the unique trace property. Preprint arXiv:1211.6395, 2012.

FACULTY OF MATHEMATICS AND INFORMATICS, UNIVERSITY OF SOFIA, BLVD. JAMES BOURCHIER 5, BG-1164 SOFIA, BULGARIA

E-mail address: nikolay.antonov.ivanov@gmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSLO, P.O.Box 1053 BLINDERN, NO-0316 OSLO, NORWAY

E-mail address: trono@math.uio.no